


Book 10

Proposition 71

When a rational and a medial (area) are added together, four irrational (straight-lines) arise (as the square-roots of the total area)—either a binomial, or a first binomial, or a major, or the square-root of a rational plus a medial (area).

Let AB be a rational (area), and CD a medial (area). I say that the square-root of area AD is either binomial, or first binomial, or major, or the square-root of a rational plus a medial (area).



For AB is either greater or less than CD . Let it, first of all, be greater. And let the rational (straight-line) EF be laid down. And let (the rectangle) EG , equal to AB , have been applied to EF , producing EH as breadth. And let (the rectangle) HI , equal to DC , have been applied to EF , producing HK as breadth. And since AB is rational, and is equal to EG , EG is thus also rational. And it has been applied to the [rational] (straight-line) EF , producing EH as breadth. EH is thus rational, and commensurable in length with EF [Prop. 10.20].

Again, since CD is medial, and is equal to HI , HI is thus also medial. And it is applied to the rational (straight-line) EF , producing HK as breadth. HK is thus rational, and incommensurable in length with EF [Prop. 10.22]. And since CD is medial, and AB rational, AB is thus incommensurable with CD . Hence, EG is also incommensurable with HI . And as EG (is) to HI , so EH is to HK [Prop. 6.1]. Thus, EH is also incommensurable in length with HK [Prop. 10.11]. And they are both rational. Thus, EH and HK are rational (straight-lines which are) commensurable in square only. EK is thus a binomial (straight-line), having been divided (into its component terms) at H [Prop. 10.36]. And since AB is greater than CD , and AB (is) equal to EG , and CD to HI , EG (is) thus also greater than HI . Thus, EH is also greater than HK [Prop. 5.14]. Therefore, the square on EH is greater than (the square on) HK either by the (square) on (some straight-line) commensurable in length with (EH), or by the (square) on (some straight-line) incommensurable (in length with EH). Let it, first of all, be greater by the (square) on (some straight-line) commensurable (in length with EH). And the greater (of the two components of EK) HE is commensurable (in length) with the (previously) laid down (straight-line) EF . EK is thus a first binomial (straight-line) [Def. 10.5]. And EF (is) rational. And if an area is contained by a rational (straight-line) and a first binomial (straight-line) then the square-root of the area is a binomial (straight-line) [Prop. 10.54]. Thus, the square-root of EI is a binomial (straight-line).

Hence the square-root of AD is also a binomial (straight-line). And, so, let the square on EH be greater than (the square on) HK by the (square) on (some straight-line) incommensurable (in length) with (EH). And the greater (of the two components of EK) EH is commensurable in length with the (previously) laid down rational (straight-line) EF . Thus, EK is a fourth binomial (straight-line) [Def. 10.8]. And EF (is) rational. And if an area is contained by a rational (straight-line) and a fourth binomial (straight-line) then the square-root of the area is the irrational (straight-line) called major [Prop. 10.57]. Thus, the square-root of area EI is a major (straight-line). Hence, the square-root of AD is also major.

And so, let AB be less than CD . Thus, EG is also less than HI . Hence, EH is also less than HK [Props. 6.1, 5.14]. And the square on HK is greater than (the square on) EH either by the (square) on (some straight-line) commensurable (in length) with (HK), or by the (square) on (some straight-line) incommensurable (in length) with (HK). Let it, first of all, be greater by the square on (some straight-line) commensurable in length with (HK). And the lesser (of the two components of EK) EH is commensurable in length with the (previously) laid down rational (straight-line) EF . Thus, EK is a second binomial (straight-line) [Def. 10.6]. And EF (is) rational. And if an area is contained by a rational (straight-line) and a second binomial (straight-line) then the square-root of the area is a first bimedial (straight-line) [Prop. 10.55]. Thus, the square-root of area EI is

a first binomial (straight-line). Hence, the square-root of AD is also a first binomial (straight-line). And so, let the square on HK be greater than (the square on) HE by the (square) on (some straight-line) incommensurable (in length) with (HK) . And the lesser (of the two components of EK) EH is commensurable (in length) with the (previously) laid down rational (straight-line) EF . Thus, EK is a fifth binomial (straight-line) [Def. 10.9]. And EF (is) rational. And if an area is contained by a rational (straight-line) and a fifth binomial (straight-line) then the square-root of the area is the square-root of a rational plus a medial (area) [Prop. 10.58]. Thus, the square-root of area EI is the square-root of a rational plus a medial (area). Hence, the square-root of area AD is also the square-root of a rational plus a medial (area).

Thus, when a rational and a medial area are added together, four irrational (straight-lines) arise (as the square-roots of the total area)—either a binomial, or a first binomial, or a major, or the square-root of a rational plus a medial (area). (Which is) the very thing it was required to show.